

A Candidate Neutron Star Associated with Galactic Center Supernova Remnant Sagittarius A East

Sangwook Park^{1,5}, Michael P. Muno², Frederick K. Baganoff³, Yoshitomo Maeda⁴, Mark Morris², George Chartas¹, Divas Sanwal¹, David N. Burrows¹, and Gordon P. Garmire¹

ABSTRACT

We present imaging and spectral studies of the supernova remnant (SNR) Sagittarius (Sgr) A East from deep observations with the *Chandra X-Ray Observatory*. The spatially-resolved spectral analysis of Sgr A East reveals the presence of a two-temperature thermal plasma ($kT \sim 1$ keV and 5 keV) near the center of the SNR. The central region is dominated by emission from highly-ionized Fe-rich ejecta. We estimate a conservative upper limit on the total Fe ejecta mass of the SNR, $M_{Fe} < 0.27 M_{\odot}$. Comparisons with standard SN nucleosynthesis models suggest that this Fe mass limit is consistent with a Type II SN explosion for the origin of Sgr A East. On the other hand, the soft X-ray emission extending toward the north of the SNR can be described by a single-temperature ($kT \sim 1.3$ keV) thermal plasma with normal chemical composition. This portion of the SNR is thus X-ray emission from the heated interstellar medium rather than the metal-rich stellar ejecta. We point out that a hard pointlike source CXOGC J174545.5–285829 (the so-called “cannonball”) at the northern edge of the SNR shows unusual X-ray characteristics among other Galactic center sources. The morphological, spectral, and temporal characteristics of this source suggest an identification as a high-velocity neutron star. Based on the suggested Type II origin for the SNR Sgr A East and the proximity between the two, we propose that CXOGC J174545.5–285829 is a high-velocity neutron star candidate, born from the core-collapse SN which also created the SNR Sgr A East.

¹Department of Astronomy and Astrophysics, Pennsylvania State University, 525 Davey Laboratory, University Park, PA. 16802

²Department of Physics and Astronomy, University of California Los Angeles, Los Angeles, CA. 90095

³Center for Space Research, Massachusetts Institute of Technology, Cambridge, MA. 02139

⁴Institute of Space and Astronautical Science, 3-1-1 Yoshinodai, Sagamihara, Kanagawa, 229-8510, Japan

⁵park@astro.psu.edu

Subject headings: Galaxy: center – supernova remnants: individual (Sagittarius A East) – X-rays: general – X-rays: ISM

1. INTRODUCTION

Sagittarius (Sgr) A East, a part of the Sgr A complex in the Galactic center, is an extended radio source with an angular size of $2'.7 \times 3'.6$ (Ekers et al. 1983). The nonthermal radio spectrum (radio spectral index $\alpha = 0.76$, where $S \propto \nu^{-\alpha}$) and the shell-like morphology suggested an identification of a supernova remnant (SNR) for Sgr A East (Jones 1974; Ekers et al. 1975; Ekers et al. 1983). The *ROSAT*, *ASCA*, and *Beppo-SAX* satellites detected diffuse X-ray emission from Sgr A East, but extensive analysis was infeasible because of the limited detector capabilities (Predehl & Trümper 1994; Koyama et al. 1996; Sidoli et al. 1999).

Recently, observations with the high angular resolution instruments aboard the *Chandra X-Ray Observatory* clearly resolved the diffuse X-ray emission interior to the Sgr A East radio shell (Maeda et al. 2002, M02 hereafter). The *Chandra* data revealed a centrally-peaked X-ray morphology of Sgr A East. The X-ray emission originates from a hot thermal plasma with an electron temperature $kT \sim 2$ keV, showing strong He α lines from highly-ionized elemental species of S, Ar, Ca, and Fe. The estimated low mass of the hot gas ($\sim 2 M_{\odot}$) and the total thermal energy ($E_{th} = 2 \times 10^{49}$ ergs) indicated a single SNR origin for Sgr A East (M02). The *Chandra* data revealed that the Fe-rich ejecta is concentrated near the center of the SNR, while other elemental species are uniformly distributed over the SNR. M02 further suggested that Sgr A East is a metal-rich, mixed-morphology (MM) SNR from a Type II SN explosion of a 13–20 $M_{\odot}$ progenitor.

Sgr A East was also observed with the *XMM-Newton Observatory* (Sakano et al. 2004, S04 hereafter). The results from the *XMM-Newton* data analysis were also consistent with a single SNR interpretation for Sgr A East. S04, however, found that a two-temperature thermal plasma ($kT \sim 1$ keV and 4 keV) was required in order to adequately describe the observed X-ray emission line features. The measured metal abundance patterns and the estimated Fe ejecta mass suggested either Type Ia or Type II origin. A weak 6.4 keV “neutral” Fe line feature was also detected with the *XMM-Newton* data. This neutral Fe line was attributed to the irradiation by Sgr A East of the molecular clouds which are also interacting with Sgr A East (S04). The discrepancies between the *Chandra* and the *XMM-Newton* data might have been caused by the low photon statistics of the *Chandra* data (S04). Effects of the relatively poor angular resolution of the *XMM-Newton* data may not be ruled out, either.

Since the early *Chandra* observations of Sgr A East presented by M02, we have performed follow-up observations of the Galactic center with *Chandra* as parts of the monitoring program of the Galactic center supermassive black hole candidate Sgr A* (Baganoff et al. 2003). As of 2002 June, the total exposure of *Chandra* observations of the Galactic center reached ~ 600 ks, representing the deepest X-ray observations of the Galactic center. Utilizing the wealth of these deep observations, we have presented the complex nature of the point-like and diffuse X-ray emission features in the Galactic center in a series of publications (Baganoff et al. 2003; Munro et al. 2003a; Munro et al. 2003c; Park et al. 2004a; Munro et al. 2004a; Munro et al. 2004b). The current deep *Chandra* data increase the photon statistics of Sgr A East by an order of magnitude, compared with that of M02. Taking advantage of the deep *Chandra* exposures, in this paper we present spatially-resolved spectral analysis of Sgr A East, which was infeasible with previous data. This approach is effective for the analysis of the ejecta material, excluding substantial contamination from the swept-up ISM, as successfully demonstrated with other Galactic SNR studies (e.g., Park et al. 2004b). The large-scale, *global* spectral analysis of Sgr A East has been performed with the previous *Chandra* and *XMM-Newton* observations, which revealed overall spectral parameters and thermal characteristics of the SNR (M02; S04). Rather than presenting such general characteristics, we concentrate in this work on the origin of Sgr A East in the context of the SN explosion types. Particularly, we draw an attention to a hard *pointlike* source detected in the northern edge of Sgr A East. The overall X-ray characteristics suggest that this source is a candidate, high-velocity neutron star (NS) associated with the SNR Sgr A East. The observations are briefly described in §2. The X-ray image and spectral analyses of the SNR Sgr A East and the candidate NS are presented in §3 and §4, respectively. We discuss key characteristics of the SNR and the NS candidate in §5. Finally, a summary is presented in §6.

2. OBSERVATIONS & DATA REDUCTION

Since 1999, the Galactic supermassive black hole candidate Sgr A* has been monitored with the Advanced CCD Imaging Spectrometer (ACIS) (Garmire et al. 2003) on board the *Chandra X-Ray Observatory* (Table 1). As of 2002 June, combining 11 *Chandra*/ACIS observations (except for ObsID 1561a, which was severely contaminated by a bright transient source within the field of view [FOV]; Munro et al. 2003b), the total exposure has reached ~ 590 ks, which is the deepest ever observation of the Galactic center region in X-rays. We utilized the data reduced by Munro et al. (2003a), as we briefly describe here. We first applied the algorithm developed by Townsley et al. (2002a) to correct the spatial and spectral degradation of the ACIS data caused by radiation damage, known as charge transfer inefficiency (CTI; Townsley et al. 2000). We then screened the data by status, grade, and

the flight timeline filter. We have also removed observation time intervals of strong flaring in the background. All individual event files were then reprojected to the tangent plane at the radio position of Sgr A* ($\text{RA}[\text{J2000}] = 17^{\text{h}} 45^{\text{m}} 40^{\text{s}}.0409$, $\text{Dec}[\text{J2000}] = -29^{\circ} 00' 28''.118$) in order to generate the composite data. The detailed descriptions of the data reduction process and the resulting broadband raw image from the composite data may be found in Munro et al. (2003a).

3. Sgr A East

3.1. X-Ray Images

The center of Sgr A East radio shell is $\sim 50''$ northeast of Sgr A* and thus the entire SNR Sgr A East was imaged within the $17' \times 17'$ FOV of the ACIS-I array during the Sgr A* monitoring observations. A “true-color” X-ray image of the Sgr A East is presented in Figure 1. Each subband image has been exposure-corrected utilizing the exposure map produced by Munro et al. (2003a), and adaptively smoothed to achieve signal-to-noise (S/N) ratio of 4 by using the *CIAO* tool *csmooth*. Sgr A* is marked near the center of the image. The bright X-ray feature around Sgr A* is emission from a massive star cluster within the inner parsec of the Galactic center. The bright diffuse X-ray emission to the immediate east of Sgr A* is the SNR Sgr A East. As reported with previous *Chandra* observations by M02, X-ray emission from Sgr A East is centrally enhanced with no apparent shell-like features (Figure 1). The enhancements of the blue emission around the center of the SNR are remarkable. As we discuss in the next section, this enhanced hard X-ray emission is primarily from the Fe He α line emission. The outskirts of the SNR show soft X-ray emission. Also evident is red, soft X-ray emission extending toward the northern side of the SNR, the so-called “plume” (M02; Baganoff et al. 2003). M02 suspected the existence of a high-velocity neutron star at the “tip” of the plume, which is physically associated with Sgr A East. With the deep exposure, a hard pointlike source CXOGC J174545.5–285829 was indeed detected there (Munro et al. 2003a). This hard pointlike source might thus be the *predicted* high-velocity neutron star candidate, which might have also produced the bow shock-like plume: hereafter, we name this NS candidate the “cannonball”. We discuss the *cannonball* in detail in § 4 and § 5.1.

3.2. Equivalent Width Images

In order to examine the overall distributions of the diffuse emission line features from Sgr A East, we construct *equivalent width* (EW) images for the detected atomic emission lines, following the method described in Park et al. (2004a). After removing all detected point sources from the broadband image (see Muno et al. [2003a] for the details of the point source detection), subband images for the line and continuum bandpasses were extracted for each spectral line of interest. These subband images were adaptively smoothed to achieve an S/N ratio of 3. The underlying continuum was calculated by logarithmically interpolating between images made from the higher and lower energy “shoulder” of each broad line. The estimated continuum flux was integrated over the selected line width and subtracted from the line emission. The continuum-subtracted line intensity was then divided by the estimated continuum on a pixel-by-pixel basis to generate the EW images for each element. In order to avoid noise in the EW maps caused by poor photon statistics near the CCD chip boundaries, we have set the EW values to zero where the estimated continuum flux is low. As discussed in Park et al. (2004a), contaminations from the cosmic X-ray background and the particle background, both of which have not been subtracted in the EW generation, are insignificant in the EW images. Although the adaptive smoothing may introduce spurious faint features, the bright, arcminute-scale features are confirmed by our spectral fits (see below) and are certainly real. Therefore, we focus on these bright features in order to qualitatively investigate the overall variation in the X-ray line emission of Sgr A East.

We present the Fe He α line ($E \sim 6.6$ keV) EW image of Sgr A East in Figure 2. The energy band selections for the line and continuum to generate this EW image are presented in Table 2. The Fe He α line EW is strongly enhanced within an $\sim 40''$ diameter region near the center of the SNR. This Fe EW distribution is consistent with the central concentration of the Fe abundance in Sgr A East as reported with the previous *Chandra* and *XMM-Newton* observations (M02; S04). The high Fe EW at the center of the SNR is thus most likely caused by the enhanced Fe abundance there, displaying the distribution of Fe-rich stellar ejecta material. (Note: The bright X-ray emission at the position of Sgr A* is featureless in the Fe EW image. This is reasonable for the continuum-dominated X-ray spectrum from Sgr A* [Baganoff et al. 2003], which further supports the utility of our EW images.) In contrast to the central Fe-rich region, the northern plume region of the SNR does not show an enhancement in Fe, which suggests that it is produced by emission from shocked ISM having metal abundances close to the average value at the Galactic center.

Unlike Fe, the EW maps of other elemental species S, Ar, and Ca do not show any significant spatial structure, and we do not explicitly present those EW images. The *uniform* EW distributions of other species are also consistent with the previously reported elemental

abundance distributions across Sgr A East.

3.3. X-Ray Spectra

We perform a spatially-resolved spectral analysis of Sgr A East utilizing high angular resolution ACIS data. For the spectral analysis of our CTI-corrected data, we use the response matrices appropriate for the spectral redistribution of the CCD, as generated by Townsley et al. (2002b). Because of various roll-angles of the individual ACIS observations and the moderate angular extension of the SNR ($\sim 3'$), Sgr A East has been detected on different ACIS-I chips depending on the roll-angles. Observations taken with deep exposures between 2002 February and 2002 June (ObsID 2943 – 3665 in Table 1), however, had similar roll-angles, which represents $\sim 90\%$ of the total exposure. Small sections of the SNR have thus been detected on a single CCD for the majority of the observations. For instance, although the bright, central Fe-rich region (i.e., “center” region in Figure 1) was detected on either the ACIS-I1 or the ACIS-I3 depending on the roll-angles, $\sim 91\%$ of the total counts in the 1 – 8 keV band were detected on the ACIS-I1. We thus use detector responses appropriate for the chip positions of the source region on the ACIS-I1 for the “center” region spectrum. We used the same method for other regional spectra. We binned the data to contain a minimum of 30 – 50 counts per channel for the spectral fittings. Even though we corrected the data for the CTI, there remain some residual effects of small gain shifts, which are usually insignificant in the spectral analysis. The gain shift at $\sim 1\%$ level may, however, confuse the spectral analysis of strong emission lines in a relatively narrow energy band such as Sgr A East spectrum containing strong Fe K lines between $E \sim 6.4$ keV – 6.9 keV. We inspected any residual gain shifts with the ACIS-I calibration data (ObsID 61184) taken close to our deepest observations (2002 May 25). We found that there were $\sim 0.1\% - 1\%$ gain shifts depending on the detector positions. We thus included a “gain fit” in the initial spectral fittings and then adjusted the gain according to the best-fit gain shift. The adjusted gain shifts are typically small ($\sim 0.6\%$), and are consistent with those from the calibration data.

The background estimates for the spectral analysis of Sgr A East are difficult because X-ray background emission in the vicinity of Sgr A East shows complex spectral and spatial structure. We were particularly cautious about the background emission features from the neutral Fe K ($E \sim 6.4$ keV) and the H-like Fe K ($E \sim 6.96$ keV) lines, because line emission from these species is prevalent across the Galactic center region (Park et al. 2004a; Munro et al. 2004a). We considered several background regions which are source-free and represent the Fe line features. We eventually used the combined background emission from

three regions which, we believe, adequately represent the “average” background spectrum for Sgr A East. The adopted background regions are presented in Figure 3. The spectral analysis and discussion in the following sections are based on this background selection.

3.3.1. Central Region

We first extracted and fitted the spectrum of the $\sim 14'' \times 24''$ region containing the bright, Fe-rich emission at the core of the SNR (“center” in Figure 1 & Figure 2). This region selection was also intended to include the central region with the highest broadband intensity, while excluding the chip-gap. Even with the small angular size of $\sim 20''$, the “center” spectrum contains significant photon statistics of ~ 7000 counts, and clearly shows the strong Fe He α line at $E \sim 6.6$ keV (Figure 4). Since M02 and S04 consistently found that thermal plasma in a collisional ionization equilibrium (CIE) can adequately describe the Sgr A East spectrum, we fitted the “center” spectrum with a CIE model (*vmekal* in XSPEC) absorbed by interstellar gas (we have also fitted the spectrum with a non-equilibrium ionization [NEI] model and the best-fit ionization timescale was indeed high [$n_e t \sim 10^{12}$ cm $^{-3}$ s], indicating a CIE plasma condition). The fitted absorption column was large ($N_H \sim 15 \times 10^{22}$ cm $^{-2}$), which was consistent with the Galactic center location of the SNR. The electron temperature ($kT \sim 2$ keV) was consistent with that reported by M02. The high Fe abundance of a few times solar was also in agreement with the results of M02 and S04.

Although a single temperature CIE model may describe the observed spectrum, we recognized a residual “hump” at $E \sim 7$ keV above the best-fit model, which appears to be a weak Fe Ly α line feature ($E \sim 6.96$ keV). In fact, the relatively poor fit ($\chi^2_\nu \sim 1.4$) was primarily caused by this excess emission. Our attempts to remove these residuals by adopting background subtraction from various regions (including the same background regions used by M02 and S04) were not successful. We thus conclude that this residual emission originates from the SNR. The detection of the Fe Ly α line is indeed statistically significant ($\sim 7\sigma$). This excess emission at $E \sim 7$ keV due to the H-like Fe line is important because an additional hot plasma component is required in order to fit this feature. The two-temperature CIE model can then improve the overall fit significantly ($\chi^2_\nu \sim 1.0$; Figure 4). In this two-component fit, we varied S, Ca, Ar, and Fe abundances while keeping them in common for both components. All other species were fixed at solar abundance because the contribution from those elemental species in the fitted energy band is negligible. We also assumed the same foreground column for both components. The results of this two-component spectral fit are presented in Table 3 and Table 4. The fitted Fe abundance is high (~ 5.8 solar) while those for S (~ 0.7), Ar (~ 1.8), and Ca (~ 1.4) are relatively low.

The electron temperatures from the best-fit model are $kT \sim 1$ keV and 5 keV, which are consistent with the results from S04. Although the electron temperature for the hard component, $kT \sim 5$ keV, appears to be unusually high for an SNR spectrum, such a temperature is required to properly describe the detected Fe Ly α line feature. This highly ionized H-like Fe line emission was not detected by the early *Chandra* observations with short exposures (M02). It was, however, suggested by the *XMM-Newton* data (S04), and is now confirmed by our deep *Chandra* observations. S04, with the *XMM-Newton* data, also reported detection of “neutral” Fe line emission ($E \sim 6.4$ keV) from Sgr A East. Although more prominent in the peripheral regions of the SNR, the neutral Fe K line was reportedly detected in the central regions of the SNR with *XMM-Newton* (photon flux $\sim 3 \times 10^{-6}$ photons cm $^{-2}$ s $^{-1}$ within 28'' radius; S04). Our deep, high angular resolution *Chandra* data do not require such a neutral Fe line feature in order to describe the observed spectrum. We place a 2σ upper limit of 1.1×10^{-6} photons cm $^{-2}$ s $^{-1}$ on the 6.4 keV Fe line photon flux.

3.3.2. “Plume” Regions

We extract the spectrum of the plumelike feature from a region between the Fe-rich center and the *cannonball* (“plume” region in Figure 1 & Figure 2). The spectrum contains ~ 5300 counts and can be fitted with a single-temperature ($kT \sim 1.3$ keV) CIE plasma model (Figure 5). In this fit, the elemental abundances for S, Ar, Ca, and Fe were varied freely, while abundances for other species were fixed at solar (Table 3 and Table 4). The Fe He α line is significantly weaker (e.g., EW ~ 800 eV) than the “center” region spectrum (e.g., EW ~ 2500 eV). The Fe abundance is also considerably lower than the central region, and is consistent with solar. These results indicate that the plume is emission primarily from a shocked ISM rather than from metal-rich ejecta. In contrast to the “center” region, hard excess emission at $E \sim 7$ keV due to the Fe Ly α line feature is not substantial in the plume region: i.e., the line detection is only marginal ($\sim 2.5\sigma$), and the line flux is an order of magnitude lower than that from the “center” region. In fact, the statistical improvement with a two-temperature plasma model is insignificant (e.g., $\chi^2_\nu \sim 1.1$ vs. 1.2). The fitted metal abundances also unchange when a two-temperature model is used. Although there appears to be a small contribution from the high temperature ($kT \sim 5$ keV) component, the overall spectrum from the plume region is thus dominated by the soft component ($kT \sim 1$ keV). Therefore, we hereafter discuss the plume spectrum based on the single-temperature model, as presented in Table 3 and Table 4.

We then extract the spectrum from a region between “center” and “plume” (“north” region in Figure 1 & Figure 2). This regional spectrum contains ~ 7600 counts. The Fe He α

line is substantially weaker than the “center” region, similar to the “plume” spectrum. On the other hand, as in the “center” region spectrum, hard excess emission at $E \gtrsim 7$ keV was noticeable for the “north” region spectrum when fitted with a single-temperature model. We thus fit this regional spectrum with a two-temperature CIE plasma model in the same way as for the “center” region (Figure 6). The fitted parameters from the two-temperature model are presented in Table 3 and Table 4. The fitted parameters are generally consistent with those from the central region. It is, however, notable that the Fe abundance in the “north” region is significantly lower than that of the “center” region. As with the other regions, we found that the 6.4 keV line is not evident in the observed “north” region spectrum, with an upper limit to the line flux of 3.7×10^{-7} photons $\text{cm}^{-2} \text{s}^{-1}$ (2σ).

4. CXOGC J174545.5–285829: *The Cannonball*

4.1. X-Ray Morphology

M02 speculated that a high-velocity neutron star associated with the SNR Sgr A East might exist at the “tip” of the plume. With deep observations, a hard pointlike source was indeed detected at the edge of the plume and designated as CXOGC J174545.5–285829 (the so-called *cannonball*) by Munro et al. (2003a). Figure 7 presents the soft and hard band images of the *cannonball*. In fact, at the edge of the plume, there are two pointlike sources separated by $\sim 3''$: the *cannonball* detected exclusively above $E \sim 3$ keV, and a soft source CXOGC J174545.2–285828 detected mostly below $E \sim 2$ keV (Figure 7). The soft source CXOGC J174545.2–285828 exhibits long-term variability (Munro et al. 2004b), and has an optical counterpart in the USNO-B1.0 catalog (#0610-0600649, $m_B = 15.27$), both of which suggest that this soft source is a foreground star.

Figure 7 reveals that the *cannonball* may not be truly pointlike, but is apparently extended: i.e., a cometary shape with a bright, pointlike “head” and a “tail” extending generally southward, which is roughly tracking back to Sgr A East. Although this source was detected $\sim 2.3'$ off-axis, the apparent extended morphology is not caused by the point spread function (PSF). In Figure 8b, the one-dimensional source intensity distribution in the north-south direction is compared to that of the off-axis PSF at the same position. The extended nature of the *cannonball* toward the south is unambiguously revealed. In Figure 8a, we present a PSF-deconvolved source image, using a maximum-likelihood algorithm (Richardson 1972; Lucy 1974). For comparison, the deconvolved image of the soft source CXOGC J174545.2–285828 (the source in the right-hand side of the image) is also presented. The *cannonball* shows a significantly extended structure which confirms the cometary morphology suggested by the raw image and the projected intensity profile. The soft source,

in contrast, appears to be resolved further into two pointlike sources. The photon statistics of the two sources, *the cannonball* and CXOGC J174545.2–285828, are similar ($\sim 1000 - 1200$ counts), and thus the morphological difference between the two is unlikely a statistical artifact. The angular extent of the *cannonball*'s tail is $\sim 3''$, which corresponds to a projected physical size of ~ 0.1 pc (Hereafter, we use a distance of 8 kpc to the Galactic center [Reid 1993]).

4.2. X-Ray Spectrum

The *cannonball* has been detected on various detector positions of the ACIS-I array because of the different roll-angles for the individual observations. These various detector positions include on-chip and chip-gap positions as well as different “rows” within the CCD. We thus use detector responses appropriate for each observation to average over all the observations for the spectral analysis of the combined data. For six observations (ObsIDs 242, 1561b, 2951, 2952, 2953, and 2954) in which the source was detected on an on-chip position, the source photons are relatively evenly spread over the ACIS-I0 (46% of the total) and the ACIS-I3 (54% of the total). We thus made a weighted sum (by detected counts) of the redistribution matrix function (RMF) files appropriate for the source locations of each CCD. The corresponding ancilliary response functions (ARF) were also averaged by weighting the counts detected in the individual observations. With the other five observations (ObsIDs 2943, 3663, 3392, 3393, and 3665), nearly all of the photons (90% of the total) were detected on the ACIS-I0. We thus use the RMF appropriate for the source detector positions on the I0 chip. The ARF files for these five observations were also averaged by weighting the counts detected in the individual observations.

The source spectrum of the *cannonball* is extracted from an $\sim 1''.8$ radius circular region centered on the source position. We tested the background spectrum separately from three regions: i.e., an annulus of inner radius $\sim 5''$ and outer radius $\sim 6''.5$ centered on the source position, a circular region of radius $\sim 11''$ in the northeast side of the source, and a $10'' \times 60''$ box region along the CCD chip-gap (for ObsIDs 2943, 3663, 3392, 3393, and 3665) to the north of the source. The choice of these background regions resulted in no significant differences in the spectral fits. In the following discussion of the spectral analysis, we assume the rectangular background region along the chip-gap (Figure 1 & Figure 3), because this background region consistently represents the source “flux variation” caused by the on- and off-chip positions. We binned the source spectrum in order to contain 20 or more counts per channel. The spectrum can best be fitted with a power law (Figure 9) and the best-fit parameters are presented in Table 5. Because of the relatively low photon statistics (~ 1000

counts), the spectrum may also be fitted with thermal plasma models. The best-fit electron temperature is remarkably high ($kT \sim 25$ keV) with extremely low metal abundances. This high temperature might not be entirely unreasonable for sources like cataclysmic variables. Although unlikely, a thermal origin of the X-ray spectrum of the *cannonball* may thus not be completely ruled out.

5. DISCUSSION

5.1. The *Cannonball*: A High-Velocity Neutron Star?

The spectral and morphological characteristics of the *cannonball* are very unusual among the sources detected within the ACIS FOV of the Galactic center: i.e., it is a bright source with a hard, continuum-dominated X-ray spectrum having no Fe line emission. This source also shows no long-term variability (Muno et al. 2004b). The *cannonball* is one of only two sources, out of ~ 2300 Galactic center X-ray sources cataloged by Muno et al. (2003a), which share such unusual X-ray characteristics. We discuss this unique source in the following sections. The observed X-ray nature of the *cannonball* suggests an identification as a high-velocity neutron star in the Galactic center. Based on its proximity to Sgr A East, we therefore propose that the *cannonball* may be a candidate neutron star born from the SN explosion which also produced the SNR, Sgr A East.

5.1.1. Spectrum and Morphology

The X-ray spectrum of the *cannonball* is best-fitted with a power law ($\Gamma \sim 1.6$), which is typical for nonthermal synchrotron emission from a neutron star’s magnetosphere. The implied high foreground absorption is consistent with that of Sgr A East, in agreement with a Galactic center location nearby the SNR. Assuming a Galactic center location, the estimated X-ray luminosity ($L_X \sim 3.1 \times 10^{33}$ ergs s $^{-1}$) is also typical for a pulsar and/or pulsar wind nebula (PWN) (e.g., Becker & Aschenbach 2002).

The deep *Chandra* images unambiguously reveal a cometary morphology of the *cannonball*. The tail is faint (only $\lesssim 100$ photons), and the estimated physical size is ~ 0.1 pc. This tail size is consistent with cometary tails detected from other Galactic high-velocity pulsars such as the Geminga pulsar (Caraveo et al. 2003), PSR B1853+01 (Petre et al. 2002), and PSR J1509–5850 (Sanwal et al. in preparation). The tail points roughly toward the south, consistent with the direction of the center of Sgr A East, given its limited photon statistics. The sky position of the source is also interesting: i.e., the *cannonball* is located, in projec-

tion, just *interior* to the radio shell boundary of Sgr A East while sitting on the “tip” of the X-ray plume. These X-ray morphologies are suggestive of a high-velocity neutron star, moving toward the north, for the origin of the *cannonball*. Considering an SNR age of $\sim 5000 - 10000$ yr (Mezger et al. 1989; Uchida et al. 1998; M02; S04) and the angular separation of $\sim 2'$ between the source and the “center” of the SNR, a velocity of $v \sim 455 - 912$ km $s^{-1}/\sin\beta$, where β is the angle between the line of sight and the actual traveling direction of the neutron star, is implied. This velocity range is in good agreement with that of typical high-velocity pulsars in the Galaxy (Cordes & Chernoff 1998).

We may make an independent estimate of the velocity of the candidate neutron star by assuming that the plume is a bow-shock produced as the high-velocity neutron star encounters the ISM. Based on the conical shape of the plume as seen by the “red” emission feature in Figure 1a, we estimate an opening angle of $\theta \sim 53^\circ$. The Mach number $Ma = [\sin(\theta/2)]^{-1}$ is thus ~ 2.2 . The velocity of the candidate neutron star is then $v = c_s Ma = \sim 880$ km s^{-1} . In this estimate, we assumed a sound speed of the ambient gas $c_s = (\gamma kT/\mu m_p)^{1/2} \sim 400$ km s^{-1} where $\gamma = 5/3$ for a monatomic, adiabatic gas, $\mu = 1$ for protons, m_p is the proton mass, and $kT \sim 1$ keV plasma in the plume region. We note that, in the velocity estimation from the Mach number, we used the projected opening angle for the bow-shock. If the inclination angle of the bow-shock was significant, the actual, de-projected opening angle could have been smaller, thus higher velocity would have been derived. Nonetheless, the velocity estimates from the two methods are in plausible agreement.

A third velocity estimate can be derived by assuming that this NS candidate produced a PWN which has reached pressure equilibrium with the ISM. Assuming that the X-ray spectrum of the *cannonball* is primarily from a PWN of the neutron star, we may derive the rotational spin-down energy loss $\dot{E}$ by using an empirical relationship between $\dot{E}$ and the power law photon index of the PWN,

$$\Gamma_{PWN} = 2.36 - 0.021 \dot{E}_{40}^{-\frac{1}{2}},$$

where $\dot{E}_{40}$ is the spin-down power in units of 10^{40} ergs s^{-1} (Gotthelf 2003). The best-fit photon index of $\Gamma_{PWN} = 1.6$ implies $\dot{E} \sim 7.6 \times 10^{36}$ ergs s^{-1} . We may assume a pressure balance between the ram pressure of the PWN, $P_{PWN} = \dot{E}/(4\pi cR^2)$ where R is the PWN radius for a spherical geometry, and the thermal pressure of the plume region, $P_{th} = 2n_e kT$. The best-fit volume emission measure (EM) from the plume region implies an electron density of $n_e \sim 7.4 f^{-\frac{1}{2}} \text{ cm}^{-3}$, where f is the X-ray emitting volume filling factor (we assumed a half-conical volume with a circular base of radius $\sim 25''$ and a height $\sim 50''$ for the “plume” region). The best-fit thermal plasma model for the plume region then implies $P_{th} (= P_{PWN}) \sim 3.1 \times 10^{-8}$ ergs cm^{-3} . (The PWN radius is accordingly estimated to be $R \sim 2.5 \times 10^{16}$ cm, which corresponds to ~ 0.2 at $d \sim 8$ kpc. This small R is consistent with

the pointlike detection of the “head” of the source by the ACIS.) P_{PWN} may be equivalent to the external ram pressure $\epsilon_k = \rho v^2$ (ρ is the mass density of the ISM derived from the plume region spectrum). The velocity is then derived to be $v \sim 550 \text{ km s}^{-1}$. We note that there were a number of assumptions in these estimates, e.g., the pressure balance $P_{PWN} = P_{th} = \epsilon_k$, the $\dot{E}$ estimate from the best-fit photon index of the PWN spectrum, and the bow-shock origin for the plume. Considering various assumptions and embedded uncertainties in the above three approaches for the velocity estimates, the agreement among all three results are remarkable. The consistency among these independent estimates of the velocity thus lends all of them additional credibility.

5.1.2. Temporal Characteristics

No evidence for long-term (observation by observation) variability of the *cannonball* was detected (Muno et al. 2004b). The proximity of this source to the chip-gap during the long series of observations makes it difficult to determine whether there is short-term (within individual observation periods) variability. With a 2–10 keV band flux of $\sim 1.9 \times 10^{-13} \text{ ergs cm}^{-2} \text{ s}^{-1}$, the *cannonball* is one of the brightest sources detected in the ACIS FOV: i.e., the 5th brightest in terms of the photon flux, and the 18th brightest in the number of counts among ~ 2300 cataloged Galactic center sources (Muno et al. 2003a). According to Muno et al. (2004b), $\sim 80\%$ of the short-term variables have a *maximum* photon flux lower than the photon flux of the *cannonball*. Also, $\sim 94\%$ of the long-term variables have a *maximum* flux lower than the *cannonball* (Muno et al. 2004b). For comparison, the nearby soft source CXOGC J174545.2–285828 with similar photon statistics was identified as a long-term variable (Muno et al. 2004b). The non-variability of the *cannonball* is thus unlikely a statistical artifact, but should correspond to the true nature of this object. The constant lightcurve with a long time-basis of $\sim 3 \text{ yr}$, particularly at the “high” flux level of this source, is remarkably unusual among the Galactic center sources. This non-variability over a few years indicates that the *cannonball* is unlikely a background AGN, and supports a neutron star identification. In fact, based on the logN-logS relation for the contribution from extragalactic sources in the Galactic center (Muno et al. 2003a), a low probability of $\sim 1.2 \times 10^{-4}$ for a detection of a background extragalactic source at the flux level of the *cannonball* within the “plume” region is implied.

The discovery of a pulsar at the position of the *cannonball* would conclusively identify this source as a neutron star. Our deep *Chandra* observations, however, used the standard 3.2 s time frame which may not be adequate for a pulsar search. We, therefore, searched for pulsations in a 40-ks archival *XMM-Newton*/EPIC observation pointed at Sgr A* (observa-

tion sequence number 0111350101). The EPIC/PN instrument has a relatively short frame time of 73.4 ms even in full-window mode, so that our pulsar search was sensitive down to periods of ~ 147 ms. Because of the poor angular resolution of the EPIC instrument, the two sources, the *cannonball* and CXOGC J174545.2–285828, are not resolved. In order to discriminate contamination from the nearby soft source CXOGC J174545.2–285828, we extract the source photons only from the hard band (3 – 10 keV). We then detect ~ 460 counts from the *cannonball*. With these *XMM-Newton* data, we do not detect significant evidence of a pulsar. We place an upper limit of $\sim 40\%$ on the pulsed-fraction between 5×10^{-5} Hz and 6.8 Hz. If the *cannonball* is a pulsar associated with Sgr A East and thus has an age of ~ 10000 yr, it may have a Vela pulsar-like periodicity of ~ 100 ms. This pulsation period is beyond the detectability of the used *XMM-Newton* time resolution. The presence of a pulsar for this neutron star candidate cannot thus be ruled out yet. Follow-up X-ray and/or radio observations with high time resolution instruments would therefore be helpful for the pulsar search for this NS candidate.

5.2. Sgr A East: Fe Ejecta Mass

The enhancement of strong Fe K line emission at the center of the SNR is remarkable. This Fe line emission is most likely from the Fe-rich stellar ejecta from the progenitor, which was heated by the reverse shock. Assuming that the Fe-rich central region represents the bulk of the total Fe mass ejected from the progenitor star, this Fe-rich material provides a useful opportunity to investigate the progenitor mass, and thus the SN explosion type of Sgr A East.

In order to estimate the total Fe ejecta mass, we first consider a simple geometry in which *all* Fe ejecta material is concentrated within the central $\sim 40''$ diameter region where the Fe EW is the highest (e.g., Fe EW $\gtrsim 1400$ eV in Figure 2). We estimate the electron density based on the measured *EM* from the spectral fit of the “center” region. We use a cylindrical geometry with an elliptical cross-section (major and minor axis radii of $12''$ and $7''$, respectively) for the X-ray emitting volume of the “center” region. A path-length of 1.55 pc was assumed, corresponding to $\sim 40''$ angular extent of the Fe enhancement at the center of the SNR. The X-ray emitting volume of $V = 1.82 f \times 10^{55} \text{ cm}^3$ is thus used for the density estimate. We, for simplicity, assume a “pure” Fe-ejecta case for the estimate of the electron density: i.e., *all* electrons originate from the ionized Fe. Considering the dominant ionization states presented by two-temperature plasma, we assume electron-Fe ion density ratio of $n_e = 24n_{Fe}$ for the He-like Fe. We also use the best-fit Fe abundance of $5.8 \times$ solar for each of the soft and the hard components. Considering the large difference in

the plasma temperatures of the two components, they are unlikely “co-spatial” (e.g., S04). We thus assume that each component occupies a separate volume and maintains a pressure equilibrium: i.e., $n_{es}T_s = n_{eh}T_h$ and $f = f_s + f_h$, where n_{es} and n_{eh} are electron densities, T_s and T_h are electron temperatures, and f_s and f_h are volume filling factors for the soft and the hard component, respectively. The best-fit electron temperatures ($kT = 1.05$ keV and 5.28 keV) correspond to $f_s = 0.48f$ and $f_h = 0.52f$. The best-fit EM s then imply Fe ion densities of $n_{Fes} \sim 0.096f^{-\frac{1}{2}} \text{ cm}^{-3}$ and $n_{Feh} \sim 0.019f^{-\frac{1}{2}} \text{ cm}^{-3}$, where n_{Fes} and n_{Feh} are Fe ion densities for the soft and the hard component, respectively. If we assume a spherical volume with a diameter of ~ 1.55 pc and with a dominant isotope ^{56}Fe , the *total* Fe mass, $M_{Fe} = 56(n_{Fes} + n_{Feh})m_pV$, is then estimated to be $M_{Fe} \sim 0.15f^{\frac{1}{2}} M_\odot$. Because of our “pure” Fe ejecta assumption, this Fe mass is an upper limit.

The standard SN nucleosynthesis models indicate that M_{Fe} is $\sim 0.5M_\odot - 0.8M_\odot$ for the Type Ia SNe (e.g., Nomoto et al. 1997a). The Type II models show a wide range of Fe mass depending on the progenitor masses: e.g., $M_{Fe} \sim 0.15M_\odot$ for a $13M_\odot - 15M_\odot$ progenitor, and $M_{Fe} \sim 0.05M_\odot - 0.08M_\odot$ for a $18M_\odot - 70M_\odot$ progenitor (e.g., Nomoto et al. 1997b). The estimated upper limit on the Fe mass for Sgr A East ($M_{Fe} \sim 0.15M_\odot$) is thus consistent with a Type II origin from a $13M_\odot - 15M_\odot$ progenitor. If our Fe mass estimates were significantly affected by the ISM contribution, the progenitor mass could be larger.

Although the total Fe mass, as estimated assuming a simple spherical geometry, provided a useful constraint on the progenitor mass, we may also entertain a contribution to the total Fe mass of the SNR from the region where Fe is marginally enhanced. Inclusion of this additional Fe mass may provide an even more conservative limit on the progenitor mass. The Fe EWs are moderately enhanced in an extended region just outside of the $\sim 40''$ Fe core (i.e., the region between the Fe EW of 800 eV and 1400 eV contours in Figure 2). This region may be represented by a “spherical shell” with an outer diameter of $\sim 1'$ and an inner diameter of $\sim 40''$. The Fe abundance in this region appears to be ~ 3 . We thus consider the contribution from this moderately Fe-rich region to the total Fe mass of the SNR. We assume that the path-length through the outer shell region (foreground and background of the Fe-core in projection) is $\sim 50\%$ of that for the inner Fe core for the adopted geometry. With the pure ejecta assumption, we consider that n_e and n_{Fe} are simply proportional to the Fe abundance ratio between the two regions (i.e., 5.8 for the inner core and 3 for the outer shell). The contribution from the shell region to the measured EM is then $\sim 10\%$. We thus use the fitted EM s from the “center” region by assuming an $\sim 90\%$ contribution from the inner Fe core (~ 1.55 pc diameter, corresponding to $\sim 40''$ angular size at 8 kpc) and an $\sim 10\%$ contribution from the outer shell (~ 0.39 pc thickness, corresponding $\sim 10''$ angular size at 8 kpc). We also assume the same fractional volume filling factors ($f_s = 0.48f$ and $f_h = 0.52f$) as above. The derived total Fe densities are $n_{Fe} \sim 0.107f^{-\frac{1}{2}} \text{ cm}^{-3}$ for the inner

Fe core and $n_{Fe} \sim 0.038 f^{-\frac{1}{2}} \text{ cm}^{-3}$ for the outer shell. We then estimate the total Fe mass from the $\sim 1'$ diameter region, $M_{Fe} \sim 0.27 f^{\frac{1}{2}} M_{\odot}$, which is $\sim 80\%$ larger than that derived merely from an $\sim 40''$ diameter spherical core. This Fe mass is again a conservative upper limit because of the pure Fe ejecta assumption. We find that this upper limit of the total Fe mass of Sgr A East is still considerably lower than that expected from Type Ia models. Based on these derived Fe mass limits, we suggest that Sgr A East was produced likely by a core-collapse SN explosion of a massive progenitor star. A core-collapse origin for the SNR Sgr A East then supports the proposed SNR-NS association between Sgr A East and the *cannonball*.

We note, however, that a Type Ia origin for Sgr A East might not be completely ruled out, yet. Considering the relatively old age of the SNR, one might speculate that a significant fraction of the Fe-rich ejecta has already been intermixed with the ambient ISM. The apparent lack of abundance enhancements from other high-Z elemental species may support this scenario. In such a case, the derived upper limits of Fe ejecta mass could be underestimates. If the amount of “missing” Fe ejecta mass was *sufficiently* large ($\sim$ several times larger than the observed Fe ejecta mass), the “true” total Fe ejecta mass might have been large enough to be consistent with a Type Ia origin for Sgr A East.

5.3. Comments on Fe Lines

The high angular resolution of the ACIS coupled with the deep exposure should be useful to investigate some controversial results between previous *Chandra* (M02) and *XMM-Newton* (S04) observations of Sgr A East: i.e., the *XMM-Newton* data reported line emission from highly-ionized H-like Fe and low-ionization state “neutral” Fe as well as the strong Fe He α line (S04), while the previous *Chandra* data showed only strong Fe He α line without H-like and neutral Fe lines (M02). These differences in the detected Fe line features resulted in discrepancies in the plasma temperatures ($kT \sim 2 \text{ keV}$ for the *Chandra* data vs. $kT \sim 1 \text{ keV}$ and 4 keV for the *XMM-Newton* data). Our data from the deep *Chandra* observations show evidence of Fe K line emission from the highly-ionized H-like Fe atoms, which confirms the results reported with the *XMM-Newton* data. The detected H-like Fe K line feature ($E \sim 6.96 \text{ keV}$) requires an extremely high temperature plasma of $kT \sim 5 \text{ keV}$ in addition to the $kT \sim 1 \text{ keV}$ component, which also confirms the results obtained with the *XMM-Newton*. We thus attribute the discrepancies between M02 and S04 on the H-like Fe line features to the poor photon statistics of the previous *Chandra* data. The hard component plasma temperature is, however, unusually high for an SNR, and the interpretation of its origin is difficult. The peculiar interstellar environments in the Galactic center, such as unusually

high magnetic fields, might be responsible for this high temperature. For instance, Munro et al. (2004a) reported the existence of very high temperature ($kT \sim 8$ keV) thermal plasma prevailing in the Galactic center regions.

On the other hand, we detect no significant evidence for the neutral Fe line emission from Sgr A East, in contrast to the results from the *XMM-Newton* observation. It is notable that the sky position of Sgr A East is on the northeastern side of Sgr A*, where the background 6.4 keV neutral Fe line emission (and other emission line features) of the Galactic center region prevails (Park et al. 2004a). The reported detection of the neutral Fe line with the *XMM-Newton* might have thus resulted from the inclusion of background diffuse emission and point sources in the spectrum of Sgr A East, neither of which could be properly accounted for, because of the poorer angular resolution of *XMM-Newton*. It is important to *avoid* confusion by the complex background line emission in the Galactic center for the spectral analysis of Sgr A East. The deep exposure and the superb angular resolution of our *Chandra* data allowed us to use small regions for the spectral analysis. For instance, our “center” region is an order of magnitude smaller in the source area than the central region used by S04, while achieving good photon statistics. We also utilized background regions which thoroughly represent *all* emission line features around Sgr A East. The detected point sources were not removed in the *XMM-Newton* data analysis (S04). Because faint point sources in the Galactic center regions are typically neutral Fe line sources (Munro et al. 2004b), the contribution from the unremoved point sources in the *XMM-Newton* data may also be responsible for the reported 6.4 keV line there (S04). In order to test these technical differences between S04 and the current work, we extracted the spectrum from a relatively large-size region ($\sim 28''$ radius) around the center of Sgr A East without removing the detected point sources. We used the *same* background region as used in S04. We then made an $\sim 4\sigma$ *detection* of the 6.4 keV line. The best-fit photon flux from the neutral Fe line was $\sim 2.5 \times 10^{-6}$ photons $\text{cm}^{-2} \text{s}^{-1}$ which was consistent with that reported by S04. We thus conclude that the previously reported 6.4 keV line emission from Sgr A East with the *XMM-Newton* data (S04) was residual contamination from the Galactic center background emission.

6. SUMMARY AND CONCLUSIONS

We presented the results from imaging and spectral analyses of the SNR, Sgr A East in the Galactic center, using deep *Chandra* observations. We confirm the central concentration of Fe-rich stellar ejecta in Sgr A East. The X-ray spectrum from the central regions of the SNR shows X-ray line emission from highly ionized He-like and H-like Fe. These Fe line features require multiple temperature thermal plasma with $kT \sim 1$ keV and 5 keV in order

to properly describe the observed spectrum. The strong Fe lines imply overabundant Fe in the center of the SNR. On the other hand, the soft X-ray emission extending to the northern side of the SNR can be described with a single temperature ($kT \sim 1.3$ keV) thermal plasma with solar abundances. This northern “plume” is thus likely X-ray emission from the shocked ISM rather than metal-rich SN ejecta.

A hard *pointlike* source (the so-called *cannonball*) detected at the northern edge of the plume, designated as CXOGC J174545.5–285829 in the Galactic center source catalog, shows remarkably unusual X-ray characteristics. The morphological, spectral, and temporal properties of this source are suggestive of an identification as a high-velocity neutron star. The estimated Fe ejecta mass of Sgr A East is consistent with a Type II SN for the origin of Sgr A East. Based on the suggested Type II origin for Sgr A East, the likely identification of the *cannonball* as a high-velocity neutron star, and the proximity between the *cannonball* and Sgr A East, we propose that these objects comprise an SNR-NS association in the Galactic center. We note, however, that follow-up pulsar searches with high time-resolution X-ray and/or radio observations at the position of the *cannonball* will be needed in order to make a conclusive identification of a neutron star for this object.

This work has been supported in parts by NASA contract NAS8-39073, NAS8-01128 for the *Chandra X-Ray Observatory*. MPM is supported by the Hubble Fellowship program through grant number HS-HF-01164.01-A.

REFERENCES

- Baganoff, F. K., Maeda, Y., Morris, M., Bautz, M. W., Brandt, W. N., Cui, W., Doty, J. P., Feigelson, E. D., Garmire, G. P., Pravdo, S. H., Ricker, G. R., & Townsley, L. K. 2003, *ApJ*, 591, 891
- Becker, A., & Aschenbach, B. 2002, in *Proc. 270th WE-Heraeus Seminar on Neutron Stars, Pulsars, and Supernova Remnants*, ed. W. Becker, H. Lesch, & J. Trümper (MPE Rep. 278; Garching: MPE), 64
- Caraveo, P. A., Bignami, G. F., De Luca, A., Mereghetti, S., Pellizzoni, A., Mignani, R. P., Tur, A., & Becker, W. 2003, *Science*, 301, 1345
- Cordes, J. M. & Chernoff, D. F. 1998, *ApJ*, 505, 315
- Ekers, R. D., Goss, W. M., & Schwarz, U. J. 1975, *A&A*, 43, 159

- Ekers, R. D., van Gorkom, J. H., Schwarz, U. J., & Goss, W. M. 1983, *A&A*, 122, 143
- Garmire, G. P., Bautz, M. W., Ford, P. G., Nousek, J. A., & Ricker, Jr., G. R. 2003, “X-Ray and Gamma-Ray Telescopes and Instruments for Astronomy”, *Proceedings of SPIE*, ed., J. E. Trümper & H. D. Tananbaum, Vol. 4851, 28
- Gotthelf, E. V. 2003, *ApJ*, 591, 361
- Jones, T. W. 1974, *A&A*, 30, 37
- Koyama, K., Maeda, Y., Sonobe, T., Takeshima, T., Tanaka, Y., & Yamauchi, S. 1996, *PASJ*, 48, 249
- Lucy, L. B. 1974, *AJ*, 79, 745
- Maeda, Y., Baganoff, F. K., Feigelson, E. D., Morris, M., Bautz, M. W., Brandt, W. M., Burrows, D. N., Doty, J. P., Garmire, G. P., Pravdo, S. H., Ricker, G. R., & Townsley, L. K. 2002, *ApJ*, 570, 671 (M02)
- Mezger, P. G., Zylka, R., Salter, C. J., Wink, J. E., Chini, R., Kreysa, E., & Tuffs, R. 1989, *A&A*, 209, 337
- Muno, M. P., Baganoff, F. K., Bautz, M. W., Brandt, W. N., Broos, P. S., Feigelson, E. D., Garmire, G. P., Morris, M. R., Ricker, G. R., & Townsley, L. K. 2003a, *ApJ*, 589, 225
- Muno, M. P., Baganoff, F. K., & Arabadjis, J. S. 2003b, *ApJ*, 598..474
- Muno, M. P., Baganoff, F. K., Bautz, M., Brandt, W. N., Garmire, G. P., & Ricker, G. R. 2003c, *ApJ*, 599, 465
- Muno, M. P., Baganoff, F. K., Bautz, M., Feigelson, E. D., Garmire, G. P., Morris, M. R., Park, S., Ricker, G. R., & Townsley, L. K. 2004a, *ApJ*, 613, 326
- Muno, M. P., Arabadjis, J. S., Baganoff, F. K., Bautz, M. W., Brandt, W. N., Broos, P. S., Feigelson, E. D., Garmire, G. P., Morris, M. R., & Ricker, G. R. 2004b, *ApJ*, 613, 1179
- Nomoto, K., Iwamoto, K., Nakasato, N., Thielemann, F. -K., Brachwitz, F., Tsujimoto, T., Kubo, Y., & Kishimoto, N. 1997a, *Nucl Phys A*, 621, 467
- Nomoto, K., Hashimoto, M., Tsujimoto, T., Thielemann, F. -K., Kishimoto, N., Kubo, Y., & Nakasato, N. 1997b, *Nucl Phys A*, 616, 79

- Park, S., Muno, M. P., Baganoff, F. K., Maeda, Y., Morris, M., Howard, C., Bautz, M. W., & Garmire, G. P. 2004a, *ApJ*, 603, 548
- Park, S., Hughes, J. P., Slane, P. O., Burrows, D. N., Roming, P. W. A., Nousek, J. A., & Garmire, G. P. 2004b, *ApJ*, 602, L33
- Petre, R., Kuntz, K. D., & Shelton, R. L. 2002, *ApJ*, 579, 404
- Predehl, P. & Trümper, J. 1994, *A&A*, 290, L29
- Reid, M. J. 1993, *ARA&A*, 31, 345
- Richardson, W. H., 1972, *J. Opt. Soc. Am.*, 62, 55
- Sakano, M., Warwick, R. S., Decourchelle, A., & Predehl, P. 2004, *MNRAS*, 350, 129 (S04)
- Sidoli, L., Mereghetti, S., Israel, G. L., Chiappetti, L., Treves, A., & Orlandini, M. 1999, *ApJ*, 525, 215
- Townsley, L. K., Broos, P. S., Garmire, G. P., & Nousek, J. A. 2000, *ApJ*, 534, L139
- Townsley, L. K., Broos, P. S., Nousek, J. A., & Garmire, G. P. 2002a, *Nuclear Instruments & Methods in Physics Research Section A*, 486, 751
- Townsley, L. K., Broos, P. S., Chartas, G., Moskalenko, E., Nousek, J. A., & Pavlov, G. G. 2002b, *Nuclear Instruments & Methods in Physics Research Section A*, 486, 716
- Uchida, K. I., Morris, M., Serabyn, E., Fong, D., & Meseroll, T. 1998, in *IAU Symp. 184, The Central Regions of Galaxy and Galaxies*, ed. Y. Sofue (Dordrecht: Kluwer), 317

Table 1. Chandra/ACIS Observation log

ObsID	Date	Exposure (ks)
0242	1999-09-21	41
1561a	2000-10-26	36
1561b	2001-07-14	14
2943	2002-05-22	35
2951	2002-02-19	12
2952	2002-03-23	12
2953	2002-04-19	12
2954	2002-05-07	12
3392	2002-05-25	167
3393	2002-05-28	158
3663	2002-05-24	38
3665	2002-06-03	90

Table 2. Energy Bands used for Generating the Iron Equivalent Width Image.

Elements	Line (eV)	Low ^a (eV)	High ^a (eV)
Fe (He α)	6510 – 6690	5000 – 5730	7470 – 8500

^aThe high and low energy bands around the selected line energies used to estimate the underlying continuum.

Table 3. Results of Spectral Fittings of Sgr A East^a

Region	N_H (10^{22} cm $^{-2}$)	kT_{soft} (keV)	kT_{hard} (keV)	EM_{soft} (10^{57} cm $^{-3}$)	EM_{hard} (10^{57} cm $^{-3}$)	χ^2/ν
Center	$18.9^{+2.3}_{-1.8}$	$1.05^{+0.19}_{-0.13}$	$5.28^{+0.66}_{-0.57}$	$7.1^{+5.5}_{-2.8}$	$0.3^{+0.1}_{-0.1}$	167.9/163
Plume	$14.7^{+1.5}_{-1.9}$	$1.32^{+0.24}_{-0.11}$	-	$1.8^{+0.6}_{-0.6}$	-	100.5/83
North	$13.0^{+1.0}_{-1.0}$	$1.00^{+0.15}_{-0.10}$	$10.9^{+19.1}_{-4.3}$	$3.7^{+1.9}_{-1.3}$	$0.05^{+0.04}_{-0.02}$	139.7/110

^aThe errors are 2σ uncertainties.

Table 4. Best-Fit Metal Abundances of Sgr A East^{a,b}

Region	S	Ar	Ca	Fe
Center	$0.69^{+0.74}_{-0.56}$	$1.76^{+0.80}_{-0.72}$	$1.37^{+0.80}_{-0.76}$	$5.81^{+1.65}_{-1.11}$
Plume	$1.91^{+0.98}_{-0.69}$	$1.20^{+0.65}_{-0.58}$	$2.50^{+0.77}_{-0.71}$	$0.96^{+0.38}_{-0.27}$
North	$1.00^{+0.31}_{-0.28}$	$1.09^{+0.44}_{-0.44}$	$1.55^{+0.63}_{-0.59}$	$2.49^{+1.36}_{-0.85}$

^aAbundances are with respect to solar.

^bThe errors are 2σ uncertainties.

Table 5. Results of Spectral Fittings of The *Cannonball*

Γ^a	N_H^a (10^{22} cm $^{-2}$)	$f_{X(2-10keV)}^a$ 10^{-13} ergs cm $^{-2}$ s $^{-1}$	$L_{X(2-10keV)}$ 10^{33} ergs s $^{-1}$	χ^2/ν
$1.59^{+0.58}_{-0.39}$	$16.5^{+3.3}_{-3.2}$	$1.9^{+3.6}_{-1.2}$	3.1	37.0/46

^aThe errors are 2σ uncertainties.




Fig. 1.— *Left*: The exposure-corrected true color image of Sgr A East from the composite data of 11 *Chandra*/ACIS observations: red represents the 1.5 – 4.5 keV band photons, green is 4.5 – 6.0 keV, and blue is 6.0 – 8.0 keV band. Each subband image has been adaptively smoothed and the detected point sources have not been removed. *Right*: The gray-scale, broadband (1.5 – 8.0 keV) ACIS image of Sgr A East. Darker gray scales are higher intensities. This broadband image is unsmoothed, and has not been exposure-corrected. The ACIS-I chip-gaps are evident: i.e., the “whiteish (or low intensity) lanes” bisecting the SNR by north-south and east-west. Some key features within the FOV and regions used for the spectral analysis are marked.




Fig. 2.— Gray-scale EW image of Fe He α line ($E \sim 6.6$ keV) from Sgr A East. Linear gray-scales (with darker gray-scales correspondign to higher intensities) range from 200 eV to 2500 eV. Three contours represent 800 eV, 1400 eV, and 2000 eV. The line and continuum images were adaptively smoothed prior to calculation of the EW. Some key features within the FOV and regions used for the spectral analysis are marked.




Fig. 3.— The *Chandra*/ACIS image of Sgr A East and its vicinity. Three circles are regions where the background emission was extracted for the SNR spectral analysis. The small rectangular box along the ACIS-I chip-gap is the background region for the X-ray spectrum of the *cannonball*. The dashed diagonal line shows the Galactic plane through Sgr A*. Darker gray scales are higher intensities. This broadband (1.5 – 8 keV) image has been rebinned by 4×4 pixels and was not exposure-corrected. Detected point sources have not been removed in this image, although the actual background spectra were extracted after removing all detected point sources.




Fig. 4.— The *Chandra*/ACIS spectrum of the Fe-rich core of Sgr A East (“center” region). The best-fit two-temperature thermal plasma model is overlaid.




Fig. 5.— The *Chandra*/ACIS spectrum of the “plume” region. The best-fit single temperature thermal plasma model is overlaid.




Fig. 6.— The *Chandra*/ACIS spectrum of the “north” region. The best-fit two-temperature thermal plasma model is overlaid.




Fig. 7.— The *Chandra*/ACIS image of the *cannonball* and CXOGC J174545.2–285828. (a) The hard band (3 – 8 keV) image. (b) The soft band (0.8 – 2 keV) image. Darker gray-scales are higher intensities.




Fig. 8.— (a) The deconvolved *Chandra*/ACIS images of the *cannonball* (left source) and CXOGC J174545.2–285828 (right source). The image has been smoothed with a Gaussian of $\sigma = 0''.25$ for the purpose of display. (b) The source intensity profiles projected in the north-south direction taken from the raw ACIS image. The histogram represents the *cannonball*, and the projected PSF at the source position is overlaid with a dashed curve.




Fig. 9.— The *Chandra*/ACIS spectrum of the *cannonball*. The upper plot is the on-chip spectrum and the lower plot is the chip-gap spectrum. The best-fit power law model is overlaid.